## Singular Integral Equations, Boundary Value Problems and the Riemann Roch Theorem

## WALTER KOPPELMAN\*

Communicated by P. R. GARABEDIAN

The Hilbert problem (often called "Riemann problem") is a boundary value probelm for analytic functions which was first formulated by Hilbert during his investigations of a set of problems mentioned by Riemann in his dissertation [19]. Of these, a certain linear boundary value problem, called the Riemann-Hilbert problem has, because of its various applications, been treated extensively in the literature recently [3, 8, 9, 15, 22]. Hilbert's work [7] and subsequently investigations [16, 11, 12] revealed a close connection between the Hilbert problem, the Riemann-Hilbert problem and the subject of one-dimensional singular integral equations. Through a study of results obtained by I. N. Vekua [22], the author was able to note a natural link between the Riemann-Hilbert problem and the Riemann-Roch theorem [9]. Our aim here is to encompass these four subjects simultaneously and to obtain various interrelations between them.

The problem considered originally by Hilbert was essentially the following: Let L be a simple closed smooth curve in the complex plane, that is, a curve which is continuously differentiable with a tangent at every point, and denote the interior of L by  $S^+$ , the exterior by  $S^-$ . It is required to find a sectionally holomorphic function  $\Phi(z)$  of finite degree at infinity, i.e.,  $\Phi(z)$  is to be holomorphic in  $S^+$  and  $S^-$  with at most a pole at infinity, such that

(1) 
$$\Phi^+(t) = G(t)\Phi^-(t) \quad \text{on} \quad L,$$

where G(t) is a given, non-vanishing Hölder continuous function on L and  $\Phi^+$ ,  $\Phi^-$  denote the boundary values of  $\Phi$  along the inner and outer edges of L, respectively.

Hilbert himself assumed more stringent conditions for G(t) and L. He trans-

<sup>\*</sup> Part of the research leading to this paper was performed under the auspices of the Atomic Energy Commission while the author was a Visiting Mathematician at Brookhaven National Laboratory. The completion of the work was supported by the United States Air Force under contract No. AF-49(638)-42, monitored by the Air Force Office of Scientific Research of the Air Research and Development Command.