

A New Class of Probability Limit Theorems

JOHN LAMPERTI*

Communicated by M. ROSENBLATT

1. Introduction. The study of the asymptotic behavior of $S_n = \xi_1 + \dots + \xi_n$, where ξ_i are independent, identically distributed, random variables with mean 0 and finite variance, is an important problem of probability theory whose solution has attained a considerable degree of perfection. The purpose of this paper is to begin to develop a theory of another rather extensive class of Markov processes along somewhat the same lines.

This vague statement can be clarified with an example: Suppose that $\{X_n\}$ is a *random walk* on the non-negative integers with the transition probabilities

$$(1.1) \quad p_{i,i+1} = \frac{1}{2} \left[1 + \frac{\alpha}{i} + o\left(\frac{1}{i}\right) \right], \quad p_{i,i-1} = 1 - p_{i,i+1}, \quad i \neq 0, \\ p_{01} = 1 - p_{00} > 0, \quad p_{ii} = 0 \quad \text{otherwise.}$$

We shall prove in § 2 that provided $\alpha > -\frac{1}{2}$, so that $\{X_i\}$ is null-recurrent or transient [6],

$$(1.2) \quad \lim_{n \rightarrow \infty} \Pr \left(\frac{X_n}{\sqrt{n}} \leq y \right) = \int_0^y \frac{2\xi^{2\alpha} e^{-\frac{1}{2}\xi^2}}{2^{\alpha+\frac{1}{2}} \Gamma(\alpha + \frac{1}{2})} d\xi, \quad y \geq 0.$$

This is our analogue of the central limit theorem giving the limit distribution of S_n/\sqrt{n} .

There are, of course, a large number of additional limit theorems for $\{S_n\}$. Many (but by no means all) of these are subsumed by the ERDOS-KAC-DONSKER *invariance principle* [3, 2] which says that, in a certain sense, the stochastic process $\{S_i/\sqrt{n}, i = 0, 1, 2, \dots\}$ converges to a Brownian motion (Wiener) process as $n \rightarrow \infty$; a large class of path functionals, such as $\max(S_1/\sqrt{n}, \dots, S_n/\sqrt{n})$, thus are shown to have limiting distributions. The greater part of the present paper is devoted to showing that a similar convergence occurs in the theory of processes like the random walks above. The limiting process is a Wiener process with reflecting barrier at the origin if $\alpha = 0$, but for other values

* This research was partially supported by the National Science Foundation. An abstract of the paper appeared in *Bull. Amer. Math. Soc.* **67** (1961), pp. 267-269.