## On the Classification of Hermitian Symmetric Spaces

## JOSEPH A. WOLF:

Communicated by S. S. Chern

- 1. Introduction. The hermitian symmetric spaces were first studied by É. Cartan [3], who classified them by means of his classification [2] of the Riemannian symmetric spaces. The work of Borel and de Siebenthal on subgroups of maximal rank in compact Lie groups [1] gives a simpler proof of Cartan's classification result. Other proofs result from H. C. Wang's work on C-spaces [7] and J. Tits' work on parabolic groups [6]. We will give a direct proof which appears to be the simplest available.
- 2. Preliminaries. In order to establish notation and terminology and to reduce our problem to a problem on compact simple Lie groups, we recall some basic notions on symmetric spaces. The reader is referred to Helgason's book [4] for details.
- 2.1. A Riemannian symmetric space is a connected Riemannian manifold M such that, given  $x \in M$ , there is a (globally defined) isometry  $s_x$  which preserves x and has differential -I on the tangent space  $M_x$ ;  $s_x$  is the symmetry to M at x. Given a complex manifold with hermitian metric, one obtains a Riemannian manifold by taking the underlying real manifold and the real part of the hermitian metric. If this Riemannian manifold is symmetric, and if the symmetries are hermitian isometries, one says that the original complex manifold with hermitian metric is a hermitian symmetric space. A hermitian symmetric space is always Kählerian.

Let  $\widetilde{M}$  be a complete simply connected Riemannian manifold. Then [5]  $\widetilde{M}$  is isometric to a product  $M_0 \times M_1 \times \cdots \times M_r$ , where  $M_0$  is a Euclidean space and the other  $M_i$  are irreducible (not Euclidean, and not locally isometric to a product of lower dimensional Riemannian manifolds.)  $\widetilde{M}$  is symmetric if and only if each of the  $M_i$  is symmetric.  $\widetilde{M}$  is the real structure of a Kähler manifold if and only if each of the  $M_i$  is the real structure of a Kähler manifold,

<sup>\*</sup> Partially supported by National Science Foundation Grant GP-812.