Construction of a Markov Process from Hitting Probabilities

FRANK KNIGHT* & STEVEN OREY*

Communicated by W. Feller

Ever since the work of Hunt [2] hitting probabilities have played a prominent part in the study of Markov processes. In [1] it was shown that for a wide class of processes the hitting probabilities characterize a Markov process with stationary transition probabilities up to a change of time scale. The question naturally arises when a class of measures $H_A(x, \cdot)$, with x ranging over points of some space K and A over subsets of K is the class of hitting distributions for some Markov process with stationary transition probabilities and state space K. When considering a Markov process X(t) and a set A of the state space let

$$T_A = \inf_{t \geq 0} \{t : X(t) \in A\}$$

be the first entrance time into A. Note that we depart from the customary definition by taking the infimum over all non-negative t and not just the positive t. The hitting probability measure $H_A(x, \cdot)$ is determined by

$$H_A(x, B) = P[X(T_A) \in B \mid X(0) = x].$$

The processes which we shall deal with all have the property that they are absorbed at a point Δ after a finite time with probability 1. The time till reaching Δ is the *lifetime* of the process.

We shall consider a compact metric space with a distinguished point Δ . We shall assume the existence of measures $H_A(x, \cdot)$ and a function g(x) satisfying certain postulates. The purpose of the paper is to show that there exists a Markov process with stationary transition probabilities having the $H_A(x, \cdot)$ as hitting probabilities and such that g(x) is the expected lifetime of the process when started at x. A precise statement, embodying also a uniqueness assertion will be found in the Theorem terminating the paper.

We are pleased to acknowledge numerous interesting conversations on the subject matter of this paper with Professor Harry Furstenberg.

^{*} This research was supported in part by the Air Force Office of Scientific Research.