A Geometric Property of the Two-dimensional Harmonic Green's Function

ZEEV NEHARI

Communicated By D. GILBARG

If D is a simply-connected domain in the xy-plane containing the point at infinity, we denote by g(x, y) the harmonic Green's function of D with singularity at infinity, i.e., $g(x, y) - \frac{1}{2} \log (x^2 + y^2)$ is harmonic in D, and g(x, y) approaches zero if the point (x, y) approaches the boundary of D from within. We shall establish the following result.

Theorem I. Let D be a simply-connected domain in the xy-plane containing the point at infinity, and denote by Γ the complement of D. If $H(\Gamma)$ is the convex hull of Γ , and if g(x, y) is the harmonic Green's function of D, then the set $H(\Gamma) - \Gamma$ is contained in the set whose points satisfy

(1)
$$0 < g(x, y) \le \frac{1}{2} \log 2.$$

The bound $\frac{1}{2} \log 2$ is the best possible. The set $H(\Gamma) - \Gamma$ will contain a point (x, y) at which $g(x, y) = \frac{1}{2} \log 2$ if, and only if, Γ is a semicircular arc.

Before we prove this result, we note that it has the following corollary.

Let Γ be a closed connected plane set of logarithmic capacity C. If $H(\Gamma)$ is the convex hull of Γ and C_1 denotes the logarithmic capacity of $H(\Gamma)$, then

$$(2) C_1 < \sqrt{2} C.$$

Indeed, C is defined by the expansion $g(x, y) = \frac{1}{2} \log (x^2 + y^2) - \log C + 0[(x^2 + y^2)^{-1}]$ for large $x^2 + y^2$, and the assertion follows from (1) and the fact that the logarithmic capacity is an increasing domain function, if it is observed that the region bounded by the level line $g(x, y) = \log r$ has the logarithmic capacity rC. However, unlike (1), (2) is not a sharp estimate, and the question as to the smallest constant K for which $C_1 \leq KC$ remains open. The best lower estimate for K we can offer is

$$K \ge 3\pi^{-2}2^{-7/3}\Gamma^{3}(\frac{1}{3}) = 1.159 \cdots$$

This research was supported by the Air Force Office of Scientific Research.