Limit Theorems Involving Capacities*

SIDNEY C. PORT

Communicated by M. Rosenblatt

1. Introduction. In this paper $\{X_n\}$ will always be an irreducible, transient, Markov chain with states in a denumerable set Ω (which, when convenient, we may label with the non-negative integers), and with n-th step transition probability matrix $P_n(x, y)$. We shall be concerned with the asymptotic behavior of the quantities $P_x(n < V_B < \infty)$ and $P_x(T_B > n)$, where $B \subset \Omega$ is a finite nonempty set, and V_B , T_B are respectively the time of first and last entrance into the set B. When the chain has an invariant measure $\pi(x)$, we shall also investigate the asymptotic behavior of the quantity

$$E_B(n) = \sum_x \pi(x) P_x(V_B \leq n).$$

Necessary notation and definitions are introduced in Sec. 2

Our principal results are as follows. In Sec. 3 we show that when the chain has an invariant measure $\pi(x)$, then if C(B) is the capacity of B, we have

(1.1)
$$e_{B}(n) = E_{B}(n) - E_{B}(n-1) \downarrow C(B), \qquad n \to \infty,$$
 and thus

(1.2)
$$E_B(n) \sim nC(B), \quad n \to \infty.$$

Moreover, $E_B(n) - nC(B)$ is nondecreasing, but the limit may either be finite or infinite. The behavior of the above quantity is closely related to that of the quantity

$$\sum_{j=0}^n P_x(j < V_B < \infty).$$

Our first task is to establish the conditions for finiteness. In Theorem 3.2 and Corollary 3.1 we show that for each finite nonempty set B and arbitrary state x, if for some state, say 0, we have

(1.3)
$$\sum_{n=0}^{\infty} \sum_{i=n+1}^{\infty} P_i(0,0) = \sum_{n=0}^{\infty} R_n(0,0) < \infty,$$

^{*} This research is sponsored by the United States Air Force under Project RAND—Contract No. AF 49(638)-700 monitored by the Directorate of Development Plans, Deputy Chief of Staff, Research and Development, Hq USAF. Views or conclusions contained in this Memorandum should not be interpreted as representing the official opinion or policy of the United States Air Force.