

Representability of Milnor's Functor k_{PL} .

ROBERT W. WEST

Communicated by A. H. WALLACE

1. Introduction. For the introduction we shall assume that the spaces under consideration are connected and have base points, and that all maps and homotopies preserve base points. $[X, B]$ denotes the set of homotopy classes $[f]$ of maps $f: X \rightarrow B$; the associated set-valued functor is denoted by $[\ , B]$. Let \mathfrak{W} be the category of finite CW complexes and maps.

Let f and g be maps from a space X to Y . Recall that f and g are said to be *weakly homotopic* if $f_* = g_*: [K, X] \rightarrow [K, Y]$ for every $K \in \mathfrak{W}$. (Here, $f_*[\varphi] = [f\varphi]$, $[\varphi] \in [K, X]$.) If in the definition of "H-space" we replace the term "homotopy" by "weak homotopy", we then get the definition of a *weak H-space*. One similarly defines the notion of *weak homotopy associativity, etc.*, of a multiplication $\mu: X \times X \rightarrow X$ on X . If X and Y are spaces with multiplications μ and ν respectively, a map $f: X \rightarrow Y$ is called a *weak homomorphism* if $f\mu$ is weakly homotopic to $\nu(f \times f)$.

A multiplication μ on B induces a multiplication (written additively) on each $[X, B]$ by $[f] + [g] = [\mu(f, g)]$ where $(f, g): X \rightarrow B \times B$ is defined by $(f, g)(x) = (f(x), g(x))$. If $f: B \rightarrow B'$ is a weak homomorphism, then $f_*: [X, B] \rightarrow [X, B']$, $X \in \mathfrak{W}$, is a homomorphism.

Milnor has developed in [5] the theory of PL-microbundles. In particular, he has defined $k_{PL}(X)$ as the set of stable equivalence classes of PL-microbundles over X . Whitney sum immediately makes $k_{PL}(X)$ an abelian monoid (it's really a group but this requires proof).

Our main result is the following.

(1.1) Theorem. *The functor k_{PL} is representable on \mathfrak{W} in the sense that there exists a countable CW complex B_{PL} such that k_{PL} is naturally equivalent to $[\ , B_{PL}]$. Addition on k_{PL} is represented by a weakly homotopy commutative and weakly homotopy associative H-space structure μ_{PL} on B_{PL} . Moreover, (B_{PL}, μ_{PL}) is unique in the sense that if (B, μ) is another H-space representing k_{PL} on \mathfrak{W} then there exists a weak homomorphism $B_{PL} \rightarrow B$ which is a weak homotopy equivalence.*

* Research was supported in part by the National Science Foundation, grant number GP-3685.