

Exotic Cobordism Theories Associated with Classical Groups

LARRY SMITH & R. E. STONG

Communicated by ANDREW WALLACE

§1. Introduction. Among the standard notions of cobordism theory is the concept of ‘cobordism of manifolds with additional structure on the normal bundle’. Briefly, one begins with a fibration

$$f : B \rightarrow BO$$

where BO is the classifying space for the stable orthogonal group O , and a (B, f) manifold is a compact differentiable manifold M^n together with a homotopy class of liftings of the normal map of $M, \nu : M \rightarrow BO$, to B . The cobordism group of closed n -dimensional (B, f) manifolds will be denoted $\Omega_n(B, f)$.

Let $f_r : B_r \rightarrow BO_r$ be the fibration induced from f by the inclusion of BO_r in BO and let $g_r : B_r \rightarrow B_{r+1}$ be the fibre map induced by the inclusion of BO_r in BO_{r+1} . The Thom space $Tf_r^*(\gamma^r)$ is formed by collapsing to a point, ∞ , all vectors of length at least one in the bundle $f_r^*(\gamma^r)$ over B_r , induced from the canonical r -plane bundle γ^r over BO_r . The fibre maps g_r induce maps

$$Tg_r : \Sigma Tf_r^*(\gamma^r) \rightarrow Tf_{r+1}^*(\gamma^{r+1})$$

where Σ denotes the suspension, defining a spectrum $\mathbf{TB} = \{Tf_r^*(\gamma^r), Tg_r\}$.

The primary result in this form of cobordism theory is the generalized Pontrjagin–Thom theorem:

The cobordism group of n -dimensional closed (B, f) manifolds is isomorphic to the n -th stable homotopy group of the spectrum \mathbf{TB} ; i.e.

$$\Omega_n(B, f) \cong \lim_{r \rightarrow \infty} \pi_{n+r} \cdot (Tf_r^*(\gamma^r), \infty).$$

Note: A full discussion of the above ideas may be found in Lashof [3].

One way to construct such cobordism theories is to be given a representation $\rho : G \rightarrow O$, G being a topological group. One then has induced a fibration on the classifying space level

$$B\rho : BG \rightarrow BO.$$