On Nonlinear Parabolic Variational Inequalities

BUI AN TON

Communicated by Hans Lewy

Let K be a closed convex subset of a real reflexive separable Banach space V with $0 \in K$ and let A be a nonlinear monotone coercive operator mapping bounded sets of $L^p(0, T; V)$ into bounded sets of $L^q(0, T; V^*)$. Brezis [2], Browder [4], Lions [8] and others have shown the existence of an element u in $L^p(0, T; V)$ with u(t) in K for almost all t, such that

$$\int_0^T (v' + Au - f, v - u) dt \ge 0$$

for all v in $L^{p}(0, T; V)$, v' in $L^{q}(0, T; V^{*})$, v(t) in K and v(0) = 0.

The main purpose of this paper is to study the regularity of the solution u when A is a nonlinear elliptic operator written in divergence form and V is a subspace of $W^{m,p}(G)$. Following Lions, we shall make a systematic use of the penalty function.

The notations and the assumptions are given in Section 1. The global regularity of u with respect to the time-variable is shown in Section 2. The results obtained, contain those of Brezis [2] and extend the local regularity results of Visik [11]. In Section 3, the case when K is a closed convex subset of both V and H with $V \subset H$ is considered. With some additional hypotheses, the regularity of u with respect to the space-variables is obtained by using an imbedding theorem of Dubinskii [6].

The regularity of solutions of linear elliptic and parabolic inequalities has been studied by Lions [8], [9], Brezis-Stampacchia [3], Lewy-Stampacchia [7].

Section 1. Let G be a bounded open subset of E^n with a boundary ∂G of class C^{∞} . Set $D_i = i^{-1}\partial/\partial x_i$; $j = 1, \dots, n$ and for each n-tuple $\alpha = (\alpha_1, \dots, \alpha_n)$ of non-negative integers, we write

$$D^{\alpha} = \prod_{j=1}^{n} D_{j}^{\alpha_{j}}$$
 with $|\alpha| = \sum_{j=1}^{n} \alpha_{j}$.

Indiana University Mathematics Journal, Vol. 22, No. 4 (1972).