

Ideal Sets and Ideals of Compact Operators on Hilbert Space

NORBERTO SALINAS

Communicated by ANDREW WALLACE

1. Introduction. In this paper we pursue the study of the two sided ideal structure of the ring $\mathcal{L}(\mathfrak{H})$ of all bounded linear operators on a separable infinite dimensional complex Hilbert space \mathfrak{H} , employing the same approach as in [1]. The fundamental tool of this approach (used implicitly throughout the present paper) is a theorem of Calkin [2, §1] which states the existence of a one to one lattice preserving correspondence between the two sided ideals of $\mathcal{L}(\mathfrak{H})$ and certain subsets of the set of all bounded sequences of non-negative real numbers called ideal sets.

Definition. A set J of bounded sequences of non-negative numbers is called an ideal set if it satisfies the following conditions:

- i) if $\{\lambda_n\} \in J$, then $\{\lambda_{\tau(n)}\} \in J$ for every permutation τ of the set Z^+ of positive integers;
- ii) if $\{\lambda_n\} \in J$ and $0 \leq \mu_n \leq \lambda_n$, for every $n \in Z^+$, then $\{\mu_n\} \in J$;
- iii) if $\{\mu_n\} \in J$ and $\{\lambda_n\} \in J$, then $\{\mu_n + \lambda_n\} \in J$.

We recall that, if \mathfrak{g} is a proper two sided ideal of $\mathcal{L}(\mathfrak{H})$, then the ideal set J , which corresponds to \mathfrak{g} under the above mentioned one to one mapping is the set of all sequences of eigenvalues (counted according to multiplicity) of all positive operators of \mathfrak{g} .

The simplest non trivial examples of ideal sets are the set C of all sequences of non-negative numbers tending to zero (corresponding to the ideal of all compact operators on \mathfrak{H}) and the subset F of C consisting of all sequences such that only finitely many terms are different from zero (corresponding to the ideal of all finite rank operators on \mathfrak{H}). C and F turn out to be the maximal and the minimal non trivial ideal sets, respectively [2]. From now on, an ideal set J will be called proper if the inclusions $F \subset J \subset C$ are proper. The classical examples of proper ideal sets are the ideal sets C_p of the p -Schatten ideals