Gaussian Sums over GL(2, Z/N) and Hecke Operators I

ROY J. FULLER

Communicated by the Editors

If R is a representation of $GL(2, \mathbf{Z}/N)$ and A is a matrix in $M(2, \mathbf{Z})$, then the Gaussian sum W(A) is defined by

$$W(A) = \sum_{g} R(g) \exp \left[2\pi i \operatorname{trace} (Ag)/N\right]$$

where g is summed over $GL(2, \mathbf{Z}/N)$. These Gaussian sums lead to a Hecke operator which acts on automorphic forms "of type R," which appeared, in a more general setting, in Shimura's paper [2], where it was suggested that the associated Dirichlet series might have an Euler product. Our goal is to calculate the "p factor" of the Euler product for the prime divisors of N. The main result is Proposition 6.

1. We begin by reviewing some definitions from [2], making the appropriate specializations. R will denote an irreducible representation of $GL(2, \mathbf{Z}/N)$ by complex matrices of size d. We always suppose that R is primitive, i.e., R is not obtained by composing a representation of $GL(2, \mathbf{Z}/M)$ with the natural homomorphism $GL(2, \mathbf{Z}/N) \to GL(2, \mathbf{Z}/M)$, where M is a proper divisor of N. Let $\mathbf{C}^+ = \{z \mid \text{Im}(z) > 0\}$ denote the upper half of the complex plane, and let \mathbf{C}^d denote the space of complex column vectors of dimension d. For each function

$$f: \mathbf{C}^+ \to \mathbf{C}^d$$
 and each $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $M(2, \mathbf{Z})$

such that det (A) > 0, we have the A-transform $f \mid A$ of weight k:

$$(f \mid A)(z) = \det (A)^{k/2} (cz + d)^{-k} f((az + b)/(cz + d)).$$

f is a cusp form of type (R, k) if the following conditions are satisfied:

- (1) The coordinate functions of f are all cusp forms of weight k with respect to the principal congruence subgroup $\Gamma_N = \{A \in \Gamma \mid A = I \pmod{N}\}$.
- (2) $(f \mid A)(z) = R(A)f(z)$ for all A in $\Gamma = SL(2, \mathbf{Z})$. S(R, k) will denote the space of cusp forms of type (R, k).