## Weakly Convergent Sequences in L∞

## SURJIT SINGH KHURANA

In [2], Tullio Zolezzi proved that in a positive measure space  $(X, \mathfrak{a}, \mu)$ , with  $X \in \mathfrak{a}$  and  $\mu(X) < \infty$ , a weakly convergent sequence in  $L_{\infty}$  is  $||\cdot||_{\mathfrak{p}}$ -convergent in  $L_{\mathfrak{p}}$  ( $1 \leq p < \infty$ ). In this note we show that this result is an immediate consequence of the fact that  $L_{\infty}$  satisfies the property D.P. strict ([1], p. 139). Also it easily follows from this that the result is also true for any p, with 0 .

**Theorem.** Let  $(X, \mathfrak{a}, \mu)$ , with  $X \mathfrak{e} \mathfrak{a}$  and  $\mu(X) < \infty$ , be a positive measure space and a sequence  $f_n \to f$  weakly in  $L_\infty$ . Then  $f_n \to f$  in  $(L_p, ||\cdot||_p)$  (0 .

Proof. Without loss of generality we take f=0. Take  $p, 1 \leq p < \infty$ . The canonical mapping  $(L_{\infty}, ||\cdot||_{\infty}) \to (L_p, ||\cdot||_p)$  is easily seen to be weakly compact and since  $L_{\infty}$  satisfies D.P. strict,  $||f_n||_p \to 0$ . In particular  $||f_n||_1 \to \infty$ . Take q, 0 < q < 1. If  $||f_n||_q \to 0$ , by taking subsequences, if necessary, we can assume that  $||f_n||_q \geq \eta > 0$ ,  $\forall n$ . Taking a subsequence if necessary and using the result that  $||f_n||_1 \to 0$ , we get  $f_n \to 0$  a.e. which implies that  $||f_n||_q \to 0$  a.e. Using the fact that  $\{f_n\}$  are uniformly bounded in  $||\cdot||_{\infty}$  (in a Banach space a weakly bounded set is norm-bounded), by dominated convergence theorem,  $||f_n||_q \to 0$ , a contradiction. This proves the result.

## REFERENCES

- A. GROTHENDIECK, Sur les applications lineares faiblement compactes d'espaces du type C(K)
  Canad. J. Math. 5 (1953), 129-173.
- 2. Tullio Zolezzi, On weak convergence in  $L_{\infty}$ , Indiana Univ. Math. J. 23 (1974), 765-766.

The University of Iowa

Date communicated: June 28, 1974