

Self Linking Numbers of Space Polygons

T. BANCHOFF

If a smooth closed space curve with nowhere vanishing curvature is pushed a small distance away from itself along its principal normal vectors to a new curve X_ϵ then the linking number of X with X_ϵ is independent of ϵ for ϵ sufficiently small and this number is called the *self-linking* number of X . This notion was first studied by Călugăreanu [3] using classical analytic techniques, and subsequently by Pohl [5] who used methods involving differential forms to establish a large number of properties of this self-linking number.

The purpose of this paper is to develop the theory of self-linking for polygons in 3-space in all of its ramifications. The proofs primarily rest upon a basic projection theorem which makes it possible to compute self-linking numbers simply and effectively by projecting to a plane and totalling numbers of apparent crossings and (signed) pairs of apparent inflection points.

The first section proves the basic theorem by deformation methods similar to those of [1]. Section 2 treats linking numbers of pairs of polygons in the same spirit and obtains a finite form of the *Gauss integral*. In section 3, we obtain integral formulas for the self-linking number of a polygon including a polygonal analogue of theorems involving total torsion of closed curves. Section 4 presents the connection with the definitions of Călugăreanu and Pohl in terms of normal variations and intersections with developable surfaces associated with the curve. Section 5 treats the behavior of self-linking under deformations, and the final section indicates how these concepts can be generalized to higher dimensional polyhedra.

James White and William Pohl have contributed helpful comments to this project and the author is grateful.

1. Definition of the self-linking number for a space polygon. Let $X = (X_0, X_1, \dots, X_{m-1})$ be a space polygon determined by a cycle of vertices in general position. We set $T_i = (X_{i+1} - X_i) / \|X_{i+1} - X_i\|$, the unit *tangent* vector in the direction of the i th edge (where all subscripts are to be reduced modulo m). Set $B_i = (T_{i-1} \times T_i) / \|T_{i-1} \times T_i\|$, the unit *binormal* vector to the oriented plane containing both edges of X at X_i .

The vectors B_i and B_{i+1} are orthogonal to T_i so $B_i \times B_{i+1}$ is a multiple of T_i . We set $f_i(X) = \pm 1$ depending on the sign of $B_i \times B_{i+1} \cdot T_i$, indicating whether