

p-adic Linear Groups with Ergodic Affine Transformations

BERTRAM M. SCHREIBER

Introduction. It has been conjectured that only compact groups can possess automorphisms which are ergodic with respect to Haar measure. This question was first raised by P. Halmos [8, page 29] and was later investigated in such papers as [12], [15], [17], [20], [24] and most recently in [2]. In this paper we are concerned with an analogous question for a wider class of mappings on a locally compact group, namely affine transformations. These mappings were introduced into ergodic theory on compact abelian groups by F. J. Hahn [6] and were studied in that context in [9] and [22] and on arbitrary compact groups in [16] and [20]. For noncompact, nondiscrete, locally compact abelian groups, it was shown independently in [17], [18] and [20] that no affine transformation can be ergodic. That the same result is valid for connected nonabelian groups was proved by R. K. Thomas [20]. On the other hand, a detailed study of the question of ergodicity of affine transformations on discrete groups was carried out by D. Jonah and the author [11]. In particular, it was shown in [11] that the only infinite discrete groups possessing ergodic affine transformations are the cyclic group \mathbb{Z} and the dihedral group D_∞ .

It follows from the result of Thomas on connected groups cited above that our problem centers around the study of affine transformations on totally disconnected groups (see Theorem 1 below). The main result of the present work (Theorem 3) is that if G is a linear algebraic group defined over a p -adic field k of characteristic zero and H is a locally compact group possessing a continuous homomorphism ρ onto a sufficiently large subgroup of the group $G(k)$ of k -rational points of G such that $\ker \rho$ is a solvable extension of a compact group, and if H possesses an ergodic affine transformation, then H is compact.

Ergodic affine transformations. Let G be a locally compact group, λ be a Haar measure on G , and T be a bicontinuous affine transformation on G , i.e., $T(x) = a\tau(x)$, $x \in G$, for some $a \in G$ and $\tau \in \text{Aut}(G)$. The mapping T is called *ergodic* if $\lambda(A)\lambda(A^c) = 0$ for any measurable subset A of G such that $T(A) = A$. More generally, if X is any Hausdorff topological