## Existence and Multiplicity Theorems for Semilinear Elliptic Equations with Nonlinear Boundary Conditions

## FRIEDHELM INKMANN

1. Notation and statement of the results. In this paper we are looking for solutions of the semilinear elliptic boundary value problem

(BP) 
$$\begin{cases} Au = f(x, u, Du) & \text{in } \Omega \\ Bu = g(x, u) & \text{on } \partial \Omega. \end{cases}$$

Here,  $\Omega$  will denote a bounded domain of class  $C^2$  in  $\mathbb{R}^N$ ,  $N \ge 2$ , i.e. the boundary  $\partial \Omega$  is a  $C^2$ -manifold of dimension N-1 such that  $\Omega$  lies locally on one side of  $\partial \Omega$ . By A we mean the differential operator

(1.1) 
$$Au = -\sum_{j,k=1}^{N} a_{jk}(x) D_{j}D_{k}u,$$

where  $a_{jk}=a_{kj}\in C(\bar\Omega)$  and A is uniformly strongly elliptic, i.e. there exists some constant  $\mu>0$  with  $\sum_{j,k=1}^N a_{jk}(x)\xi^j\xi^k\geq \mu|\xi|^2$  for all  $x\in\bar\Omega$  and  $\xi\in\mathbf R^N$ .

The boundary operator B is defined by

$$Bu = \frac{\partial u}{\partial \beta},$$

where  $\beta \in C^1(\partial\Omega, \mathbf{R}^N)$  is an outward pointing, nowhere tangent vector field on  $\partial\Omega$ .

For the nonlinearities f and g we suppose throughout this paper

(f1) 
$$\begin{cases} f: \bar{\Omega} \times \mathbf{R} \times \mathbf{R}^N \to \mathbf{R} \text{ is continous and } \partial f/\partial \xi, \, \partial f/\partial \eta \\ \text{exist and are continous, too. } ((x, \xi, \eta) \text{ denotes a} \\ \text{generic point of } \bar{\Omega} \times \mathbf{R} \times \mathbf{R}^N.) \end{cases}$$

(f2) 
$$\begin{cases} \text{There is an increasing function } c: \mathbf{R}_+ \to \mathbf{R}_+ \text{ such that} \\ |f(x,\xi,\eta)| \le c(|\xi|) \ (1+|\eta|^2) \text{ for all } (x,\xi,\eta) \in \bar{\Omega} \times \mathbf{R} \times \mathbf{R}^N. \end{cases}$$

(g) 
$$g: \partial \Omega \times \mathbf{R} \to \mathbf{R}$$
 is locally Lipschitz continuous.