

Helical Submanifolds in Euclidean Spaces

YI HONG

1. Introduction. Let M^n be a connected n -dimensional ($n > 1$) submanifold of an m -dimensional Euclidean space E^m . If every geodesic of M , considered as a curve in the ambient space, has all Frenet curvatures constant and these curvatures are independent of the chosen geodesic, then M^n is called a helical submanifold of E^m [8]. A helical submanifold is said to be of order d if each geodesic is properly contained in a d -plane of E^m . Helical submanifolds were investigated in [3], [4], [5], [7], [8], [9]. In [3], [4] Chen and Verheyen classified helical surfaces in E^m ($m = 3, 4, 5$). They showed that a helical surface in E^m ($m = 3, 4, 5$) is a 2-plane or the second standard immersion of a 2-sphere (i.e., a 2-sphere or a Veronese surface). In [5], Houh and the author proved that there is no surface M helically immersed into E^m of order 3.

Several years ago, B. Y. Chen conjectured that any helical submanifold of E^m is either spherical or an open portion of an n -plane. In this article, we show that any compact helical submanifold in E^m with a transitive isometry group is spherical. We also show that a helical surface of order 4 must be of positive constant curvature, and classify all helical immersions of compact rank one symmetric spaces.

I wish to express my gratitude to my advisor Professor C. S. Houh for his guidance. I also would like to thank professors B. Y. Chen, J. Ludden and R. Howard for their advice which improved the results. Special thanks go to the referee who gave many suggestions which improved the proofs of several theorems.

2. Notations and basic formulas. Assume that M is an n -dimensional submanifold immersed into E^m . Let $\tilde{\nabla}$ and ∇ be the covariant derivatives of M and E^m , respectively. For any two vector fields X, Y tangent to M , the second fundamental form h is given by

$$h(X, Y) = \tilde{\nabla}_X Y - \nabla_X Y.$$

For any vector field normal to M , we put

$$\tilde{\nabla}_X \xi = -A_\xi X + \nabla_X^\perp \xi,$$

where $-A_\xi X$ and $\nabla_X^\perp \xi$ denote the tangential and normal components of $\tilde{\nabla}_X \xi$ respectively. Then ∇^\perp is called the normal connection of the normal bundle $N(M)$.