## Erratum: Representation and Duality in Weighted Spaces of Analytic Functions, Vol. 34 (1985), 319–335

## DANIEL H. LUECKING

It has been pointed out by K. Stroethoff and others that there is a flaw in the proof of the second assertion in Corollary (3.6) of [1]. If  $\omega$  satisfies  $C_q$  (defined with respect to pseudohyperbolic disks of radius r), then it is trivial that  $\omega$  will satisfy  $C_q$  using disks of smaller radius  $r_0 < r$ . What needs to be done in order to verify that the class  $C_q$  is independent of choice of r is to show that  $\omega(E(a,r_0)) \leq C\omega(E(a,r))$  whenever  $r_0 > r$  and  $\omega$  satisfies  $C_q$  with respect to r. Inequality (3.7) certainly has the larger r on the left but the  $C_q$  assumption is there based on the larger r rather than the smaller. In essence we must be able to prove (3.7) from a weaker assumption. Since Corollary (3.8) relies on Corollary (3.6), its proof is also flawed.

We can correct the flaws in both proofs by proving (3.8) using only the correct portion of (3.6) and using (3.8) to finish Corollary (3.6). The details follow.

Suppose  $\omega$  satisfies  $C_q$  for some  $r \in (0,1)$ . Using (3.7) with  $r_0 = r_0/2$  (the correct portion of Corollary (3.6)), we have  $\omega\big(E(a,r)\big) \leq C\omega\big(E(a,r/2)\big)$ . The proof of Corollary (3.8) as given suffices to establish that  $\widetilde{\omega}(z) \leq C\widetilde{\omega}(\zeta)$  provided  $\rho(z,\zeta) < r/2$ . If  $\rho(z,\zeta) < s < 1$ , then there exist  $z_0,z_1,\ldots,z_n$ , with n depending only on s, such that  $z_0 = z$ ,  $z_n = \zeta$ , and  $\rho(z_k,z_{k-1}) < r/2$ ,  $k = 1,2,\ldots,n$ . Then  $\widetilde{\omega}(z) \leq C^n\widetilde{\omega}(\zeta)$  and Corollary (3.8) is established. Now assume again that  $\omega$  satisfies  $C_q$  for  $r \in (0,1)$ . Then  $\omega' = \omega^{-q'/q}$  satisfies  $C_{q'}$  for this same r. Let  $r_0 > r$ . We need to show that  $\omega\big(E(a,r_0)\big) < C\omega\big(E(a,r)\big)$  and a similar inequality for  $\omega'$ . The set  $E(a,r_0)$  can be covered with  $M = M(r,r_0)$  sets  $E(a_j,r)$ , where  $a_j \in E(a,r_0)$  (so that  $\rho(a_j,a) < r_0$ ). Then

$$\omega(E(a,r_0)) \leq \sum_{j=1}^{M} \omega(E(a_j,r))$$
$$= \sum_{j=1}^{M} \widetilde{\omega}(a_j) m(E(a_j,r)) \leq$$