Pointwise A Priori Bounds for Strongly Coupled Semilinear Parabolic Systems

REINHARD REDLINGER

Dedicated to Professor Wolfgang Walter on the occasion of his sixtieth birthday

§1. Introduction. In this paper we consider semilinear parabolic boundary value problems of the form

$$\frac{\partial u}{\partial t} + L(t,x)u = f(t,x,u,Du,\dots,D^{2m-1}u)$$
 in $J \times \Omega$,

(1) $B(t,x)u = 0 \quad \text{on } J \times \partial \Omega.$

where Ω is a bounded domain in \mathbb{R}^n , J=(0,T) with $T\leq\infty$, and L is a strongly coupled, uniformly elliptic operator of order 2m acting on N-vectorvalued functions u. By B we denote a homogeneous, possibly time-dependent linear boundary operator for m = 1, and the Dirichlet boundary operator B_D for m > 1. It is the purpose of this paper to establish pointwise a priori bounds for solutions of (1). The same problem in case m=1, $B=B_D$ and f=f(t,x,u)has been previously treated by Cosner [6, 7] and Alikakos [2]. (In [2] and [6] only time-independent operators L and nonlinearities f = f(x, u) are considered.) In these articles the a priori estimates are obtained by first getting a bound on the $W^{1,2}$ -norm of the solution via a Ljapunov function and then using regularity results to improve the bound to a pointwise one. We use the same approach, but with a Ljapunov function that only yields a bound in L^2 . This makes it possible to considerably weaken the structure conditions imposed on the system, but necessitates a more restrictive condition on the growth of the nonlinearity. Thus, in contrast to [2, 6, 7] we do not suppose that f has gradient structure or that L is self-adjoint.

The present paper makes essential use of results and methods developed by Amann in [4, esp. §§6, 7]. In this paper, global existence theorems for classical solutions of the boundary value problem (1) are established. The structure conditions on L and f in [4, §7] are analogous to those imposed by Cosner and Alikakos.