## The Fredholm Structure of an n-Multicyclic Operator

## DOMINGO A. HERRERO

§1. Introduction. In a series of articles, the author has analyzed the spectral structure of multicyclic and rationally multicyclic operators [13] [14] [16] (see also the joint article [17], in cooperation with L. Rodman; a large part of these results have been included in Chapters 11 and 13 of the monograph [3] [15]). A (bounded linear) operator T, acting on a complex separable infinite—dimensional Banach space  $\mathcal{X}$ , is called n-multicyclic if

(\*) 
$$\mathcal{X} = \bigvee \big\{ p_1(T)y_1, p_2(T)y_2, \dots, p_n(T)y_n : \\ p_1, p_2, \dots, p_n \text{ run over all polynomials} \big\}$$

for a suitable n-tuple of vectors  $(y_1, y_2, \ldots, y_n) \in \mathcal{X}^{(n)}$ , the direct sum of n copies of  $\mathcal{X}$ . Here  $\bigvee$  denotes "the closed linear span of", and  $(y_1, y_2, \ldots, y_n)$  is called a multicyclic n-tuple for T. The operator T is called rationally n-multicyclic if there exists  $(y_1, y_2, \ldots, y_n)$  in  $\mathcal{X}^{(n)}$  (a rationally multicyclic n-tuple for T) such that  $(\star)$  is satisfied when  $p_1, p_2, \ldots, p_n$  run over all rational functions with poles outside the spectrum  $\sigma(T)$ , of T.

Recall that an operator T is semi–Fredholm if ran T is closed, and either  $\ker T$  or  $\frac{\chi}{\operatorname{ran} T}$  is finite–dimensional. In this case, the semi–Fredholm index of T is defined by

$$\operatorname{ind} T = \dim \ker T - \dim \frac{\chi}{\operatorname{ran} T} = \dim \ker T - \dim \ker T^* \,.$$

If ind T is finite, then T is called a Fredholm operator. The open set

$$\rho_{s-F}(T) = \left\{ \lambda \in \mathbb{C} : \lambda - T \text{ is semi-Fredholm} \right\}$$

is the semi-Fredholm domain of T. (The reader is referred to [18] for the properties of semi-Fredholm operators.) If T is either n-multicyclic, or rationally n-multicyclic, then  $\dim \ker(\lambda - T)^*$  cannot exceed n, and therefore  $\operatorname{ind}(\lambda - T) \geq -n$  for all  $\lambda \in \rho_{s-F}(T)$  [13].