## On Elliptic Partial Differential Equations and Uniqueness Theorems for Closed Surfaces<sup>1</sup>

## PHILIP HARTMAN

Communicated by S. S. Chern

1. Theorems in differential geometry. This paper deals with the conditions of differentiability for the validity of several theorems concerning closed, 2-dimensional surfaces in Euclidean 3-space. The emphasis in this paper is on the portion of the proofs involving elliptic partial differential equations. In particular, most of the paper deals with the behavior of a solution of a linear, elliptic partial differential equation in the vicinity of a zero.

Consider the following uniqueness statement:

(A) Let  $F(H, K, \lambda, \mu, \nu)$  be defined on the set H > 0,  $H^2 \ge K > 0$ ,  $\lambda^2 + \mu^2 + \nu^2 = 1$  and satisfy

(1) 
$$\frac{\partial F}{\partial k_1} \frac{\partial F}{\partial k_2} > 0, \text{ where } 2H = k_1 + k_2 \text{ and } K = k_1 k_2.$$

Then, up to translations, there is at most one closed surface S of positive Gaussian curvature such that the mean and Gaussian curvatures, H and K, at the point of S where the inward unit normal of S is  $(\lambda, \mu, \nu)$  satisfy the relation F = 0.

This assertion is due to A. D. Alexandroff when F and S are analytic. It was proved by Pogoreloff under the conditions that F and S are sufficiently differentiable. His assumptions have been lightened successively by Wintner and myself ([9], where earlier references are given) and by Bers [2]. In both [2] and [9], the assumptions imply that S is of class  $C^3$  (in fact, S has local parametrizations in terms of functions possessing Hölder continuous third derivatives). The question was raised in [9] whether (A) holds under hypotheses which do not imply that S is of class  $C^3$ . This question will be answered below in the affirmative.<sup>2</sup>

<sup>&</sup>lt;sup>1</sup> The preparation of this paper was sponsored by the Office of Naval Research, U. S. Navy. Reproduction in whole or in part is permitted for any purpose of the United States Government. 
<sup>2</sup>Added March 17, 1958. Since this paper has been submitted, additional results of ALEXANDROFF [Vestnik, Leningrad University, 1956, No. 19, pp. 1–17] and POGORELOFF