Spectral Representation of Certain Semi-Groups of Operators

CASSIUS IONESCU TULCEA

Communicated by E. HILLE

Contents

- 1. The semi-group algebra. Representations.
- 2. Characters.
- 3. Hermitian spectral families.
- 4. Semi-groups of bounded normal operators.
- 5. Semi-groups of bounded scalar operators.
- 6. Direct sums of hermitian spectral families.
- 7. Semi-groups of unbounded normal operators.

Summary. In section 1 some definitions and results are given which will be used in the rest of the paper. The main results are proved in sections 4 and 7. In section 2 the characters of a certain class of semi-groups are studied. Section 5 deals with semi-groups of bounded scalar operators. In sections 3 and 6 a series of propositions on hermitian spectral families are given.¹

- 1. The semi-group algebra. Representations. Let G be a locally compact group, S a locally compact subspace of G and m the restriction of a left invariant Haar measure to S. Suppose
 - (1) $st \in S$ if $s \in S$, $t \in S$,
 - (2) m(U) > 0 for every non-void set $U \subset S$, open in S.

Let M(S) be the vector space of complex Radon measures defined on S, with compact support; for every measure μ let $S(\mu)$ be its support. For $\mu \in M(S)$, $\nu \in M(S)$ we can define the product $\mu\nu$ by means of the relations

$$\int f(s) \ d\mu\nu(s) = \iint f(st) \ d\mu(s) \ d\nu(t) \quad \text{for} \quad f \in \Re(S);$$

¹ The author is indebted to S. Kakutani for fruitful remarks concerning sections 2 and 5 and to D. A. Edwards for reading the manuscript and for discussions concerning the last section.