On Yilmaz' New Approach to General Relativity

BANESH HOFFMANN*

I. The Theorem of Yilmaz. YILMAZ¹ has presented a theory of gravitation in which he regards the metrical tensor $g_{\mu\nu}$ as a functional of a scalar function φ . His basic equations are

$$R_{\mu}^{"} - \frac{1}{2} \delta_{\mu}^{"} R = 8\pi T_{\mu}^{"}$$

and

(2)
$$g^{\mu\nu}\varphi_{,\mu\nu} \equiv (-g)^{-\frac{1}{2}} \{g^{\mu\nu}(-g)^{\frac{1}{2}}\varphi_{/\mu}\}_{/\nu} = 0$$

where the comma denotes the covariant derivative with respect to the metric $g_{\mu\nu}$, the solidus denotes the partial derivative, and the tensor T°_{μ} is defined by

(3)
$$T^{"}_{\mu} = \frac{1}{8\pi} \left(2g^{"}_{\varphi/\mu}\varphi_{/\lambda} - \delta^{"}_{\mu}g^{\lambda\tau}\varphi_{/\lambda}\varphi_{/\tau} \right),$$

this being the usual energy-momentum tensor for the scalar field $\varphi/(8\pi)^{\frac{1}{2}}$. He exhibits a static field based on the following remarkable theorem that he discovered.

Take the line element to have the special form

(4)
$$ds^2 = g_{\mu\nu} dx^{\mu} dx^{\nu} = e^{-2\varphi} dt^2 - e^{2\varphi} (dx^2 + dy^2 + dz^2)$$

with φ independent of time, and note that, for this line element, the field equation (2) reduces to the "flat" Newtonian equation

(5)
$$\nabla^2 \varphi \equiv \frac{\partial^2 \varphi}{\partial x^2} + \frac{\partial^2 \varphi}{\partial y^2} + \frac{\partial^2 \varphi}{\partial z^2} = 0.$$

Now compute the Einstein tensor $R^*_{\mu} - \frac{1}{2} \delta^*_{\mu} R$ for the metric in (4), and also compute the tensor T^*_{μ} defined in (3) using this metric. Then the two expressions will be identical—except for one discrepancy in the 4–4 components. We have, in fact, for this special line element,

^{*}Part of this work was done while the author was on leave at the University of London, King's College, and was assisted by WRIGHT A.D.C., A.R.D.C., U.S.A.F. through its European Office.

¹H. YILMAZ, Phys. Rev. 111, 1417 (1958).